

National Aeronautics and
Space Administration
Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California


Atmospheric Infrared Sounder



Cloud Properties from AIRS: Cirrus, and initial comparisons to CloudSat

by
Brian Kahn

**AIRS Science Team Meeting
Greenbelt, MD
September 28th, 2006**



National Aeronautics and
Space Administration


Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California



Atmospheric Infrared Sounder

Outline

- **Fast model retrievals of thin cirrus cloud optical depth (τ_{VIS}) and effective diameter (D_e)**
 - An illustrative AIRS granule
 - Global oceanic $\pm 30^\circ$ latitude statistics for September 6th, 2002
 - Version 4 versus 5
 - Comparison to MODIS Collection 4 operational τ_{VIS} and D_e
- **Initial comparisons of AIRS operational cloud fields to CloudSat**
 - A few vertical x-sections of AIRS cloud top height (Z_C) and effective cloud fraction (f) for both layers
- **Ongoing and future work**



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder



The fast retrieval approach

- **Combine OPTRAN clear-sky radiances with a thin cirrus parameterization**
 - Clear + cloudy sky used to fit AIRS radiances [*Yue et al., J. Atmos. Sci.* 2006]
- **Cirrus represented by series of D_e distributions using assumed habit models**
 - Here we use *Baum et al.* [2005] models
- **Minimize χ^2 of observed and simulated AIRS radiances: best τ_{VIS} and D_e**
- **Valid for ice clouds with:**
 - $\tau_{\text{VIS}} \leq 1.0$ (retrieved using thermal IR, but converted to visible optical depth)
 - $10 \mu\text{m} \leq D_e \leq 120 \mu\text{m}$
 - Single-layered cloud (according to AIRS)
- **Explore relationships between T_C , D_e , τ_{VIS} , etc.**
 - An example granule
 - Global oceans $\pm 30^\circ$ latitude




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California


Atmospheric Infrared Sounder



An Illustrative Granule: September 6th, 2002, granule 10



$T_b @ 960 \text{ cm}^{-1} \text{ (K)}$



Upper CTP (hPa)




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California


Atmospheric Infrared Sounder



Optical depth (left) and effective diameter (right) retrievals



τ_{VIS}



D_e (μm)




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California


Atmospheric Infrared Sounder



Minimum χ^2 versus optical depth and effective diameter



Minimum χ^2



Minimum χ^2
vs. τ_{VIS}

Minimum χ^2
vs. D_e (μm)



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder



In situ r_e - T_C Relationships




FIG. 1. Estimates of the ratio of ice water content to extinction from earlier studies.

- IWC/σ used as a proxy for r_e
- Increasing r_e with T_C
- Significant differences from different in situ campaigns and modeling studies
- Use to compare against AIRS




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California


Atmospheric Infrared Sounder



TOVS Path B D_e - T_C Relationships



- TOVS Path-B + ECMWF re-analysis (left)
- Generally increasing D_e with T_C
 - However, not necessarily true of thinnest cloud in tropics!
 - Same pattern seen with AIRS retrievals (below)
- TOVS $D_e > \text{AIRS } D_e$



Stubenrauch et al., Atmos. Res. [2004]




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California



Atmospheric Infrared Sounder

AIRS D_e vs. T_C for Varying Optical Depth (V4)



Middle curve: *Garrett et al. [2003]* T_C vs. r_e ; 1- σ curves on either side




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California



Atmospheric Infrared Sounder

AIRS D_e vs. T_C for Varying Optical Depth (V4, alt channel list)



Middle curve: *Garrett et al. [2003]* T_C vs. r_e ; 1- σ curves on either side




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California



Atmospheric Infrared Sounder

AIRS D_e vs. T_C for Varying Optical Depth (V5)



Middle curve: *Garrett et al. [2003]* T_C vs. r_e ; 1- σ curves on either side




National Aeronautics and
Space Administration


Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California



Atmospheric Infrared Sounder

AIRS D_e vs. T_C for Varying T_C Error (Version 5)





National Aeronautics and
Space Administration


Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California



Atmospheric Infrared Sounder

Summary of Thin Cirrus Tendencies

- Two D_e modes (10–15; 25–50 μm) at lowest τ_{VIS}
- Smallest mode fades away at larger τ_{VIS} ; larger mode increases in D_e with τ_{VIS}
- Small D_e mode composed of mixture of small and large T_C error: probably a real feature, but less robust than larger D_e mode
 - Version 4 and 5 comparisons + channel sensitivity highlight tenuous nature of small D_e mode
- Variable correlation of τ_{VIS} and T_C across various ranges of D_e



National Aeronautics and
Space Administration



Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California


Atmospheric Infrared Sounder



MODIS and AIRS Optical Depth

- Collocated, single-layer (according to AIRS), ECF ≤ 0.4 only
- For Granule 11 on September 6th, 2002 only
- AIRS clouds optically thinner and colder than MODIS
- Most MODIS retrievals warmer than 280 K: water clouds





National Aeronautics and
Space Administration



Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder



MODIS and AIRS Particle Size

- AIRS and MODIS see two particle modes, but AIRS with $D_e = 10\text{--}15\text{ }\mu\text{m}$
- MODIS $r_e = f(T_C)$ near $20\text{ }\mu\text{m}$
- Also hint of third large mode in AIRS for similar particle size as MODIS





National Aeronautics and
Space Administration


Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder



CloudSat Cloud Mask + AIRS 2-layer Clouds

CloudSat Track



AIRS CLOUD HEIGHT

AIRS CLOUD AMOUNT




Small



Large

CloudSat Cloud Mask Confidence Interval; 20 or more is confident cloud



Isolated tropical convection



National Aeronautics and
Space Administration


Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder





CloudSat Cloud Mask + AIRS 2-layer Clouds

CloudSat Track



AIRS CLOUD HEIGHT

AIRS CLOUD AMOUNT



Southern ocean cold air Cu and St



National Aeronautics and
Space Administration


Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder





CloudSat Cloud Mask + AIRS 2-layer Clouds

CloudSat Track



AIRS CLOUD HEIGHT

AIRS CLOUD AMOUNT



Subtropical low and midlevel cloudiness



National Aeronautics and
Space Administration


Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California



Atmospheric Infrared Sounder


CloudSat Cloud Mask + AIRS 2-layer Clouds

CloudSat Track



AIRS CLOUD HEIGHT


AIRS CLOUD AMOUNT



Small



Large



Winter storm in southern hemisphere



National Aeronautics and
Space Administration


Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California



Atmospheric Infrared Sounder



CloudSat Cloud Mask + AIRS 2-layer Clouds

CloudSat Track




AIRS CLOUD HEIGHT

AIRS CLOUD AMOUNT



Scattered Cbs and Ci



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder



Summary and Conclusions

- **Thin cirrus retrievals reveal bi-modal particle size behavior; not inconsistent with tropical cirrus clouds**
 - Uniform, tenuous, laminar, small particle size $> 15 \mu\text{m}$
 - Thicker, structured, larger particle size $< 15 \mu\text{m}$ (e.g., *Comstock et al., J. Geophys. Res.* [2002])
 - Need to explore why bi-modal in AIRS, not as much as *in situ* data
- **Matched-up MODIS cirrus retrievals show systematic differences to AIRS**
 - Larger optical depth and particle size at lower altitudes
 - AIRS picks up mode near $10\text{-}15 \mu\text{m}$, MODIS does not
- **CloudSat comparisons reveal usefulness of 2-layer AIRS clouds**
 - CALIPSO will better determine validity of small particle mode in AIRS



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California



Atmospheric Infrared Sounder

Current and Future Work

- **Retrieve thin cirrus properties for longer time periods**
 - Differences for land/ocean, day/night, V4 and V5, channel selection, etc.
 - Seasonal, regional, latitudinal dependencies?
 - Further comparisons to MODIS: IR versus VIS/near-IR retrieval methods
- **Collaboration with UCLA, modification to include scattering**
 - How do results change with different sampling of cirrus clouds (e.g., thin vs. thin+thick)?
 - Further comparisons to *in situ*, satellite data (e.g., MODIS)
 - Cirrus retrievals in V6 as suggested by L. Strow?
- **Further comparisons with CloudSat as more data is released**
 - Global statistics, IWC/LWC profiles, cloud typing
 - Fold in CALIPSO data
 - Fold in MODIS to understand heterogeneity, cloud type sensitivity, etc.